



Brian E. Whipker¹



Danielle Clade¹



Patrick Veazie¹

Volume 12 Number 36 October 2023

Mimics: Mexican Sunflower

Upon inspection of some problematic Mexican sunflower plants with lower leaf interveinal chlorosis and purplish-black coloration, the initial diagnosis pointed towards a direction, while the data from the leaf tissue and substrate samples supported a different conclusion. Thus this e-GRO Alert points out the importance of supportive data and knowing possible problems when diagnosing plant disorders.

Mexican sunflower or red sunflower (*Tithonia rotundifolia*) is a tall growing background plant that attracts bees, butterflies, and birds. During a grower visit, we were asked to look at a set of plants to help diagnose a case of lower leaf interveinal chlorosis (Figs. 1 & 2) and purplish-black coloration (Figs. 3 & 4). The symptomatic plants were the remainders from a landscape order that the grower fulfilled. As a result, they were mature, in bloom, and fertilization was sporadic.



Figure 1. Intermediate symptoms of interveinal chlorosis on the lower foliage. Normally this is associated with low magnesium (Mg) concentrations, but in this case, potassium (K) was the limited element. (Photo: Brian Whipker)

In North Carolina, our irrigation water is normally low in magnesium (Mg) because we do not have limestone bedrock to supply Mg. Therefore, when one observes lower leaf interveinal chlorosis, there is a very high probability that the cause is due to insufficient Mg. It is also common that late-season crops are more likely to develop a Mg deficiency after any Mg from the dolomitic limestone, which is added to adjust the substrate pH, has leached out.

www.e-gro.org

¹NC State University, Dept. of Hort. Science
bwhipker@ncsu.edu

2023 Sponsors



American
Floral
Endowment

Research
Internships
Scholarships
Education

Funding the Future of Floriculture

Ball®

fine



GRIFFIN
GREENHOUSE & NURSERY SUPPLIES



P.L. LIGHT SYSTEMS
THE LIGHTING KNOWLEDGE COMPANY

Reprint with permission from the author(s) of this e-GRO Alert.

Advanced Mg deficiency can also develop into a purplish-black coloration on the lower leaves. This adds uncertainty because low substrate pH levels (<~5.0) result in excess iron (Fe) and manganese (Mn) accumulations, which also results in purplish-black lower leaves. This situation also tends to occur late in the season as the dolomitic limestone leaches out.

With the two primary suspects identified, with an estimated 90% confidence level, the grower was told about the initial diagnosis. To confirm the diagnosis, tissue and substrate tests were necessary. The tests came back with a different conclusion.

Leaf symptomology mimics do occur, which is why visual diagnosis should always be confirmed with testing. This is especially true if the diagnosticians are not familiar with the plant species. None of us had grown Mexican sunflowers before and we did not know what to expect.

Substrate Test Results. The substrate test results primarily reported overall low fertility levels [electrical conductivity (EC) at 0.33 mS/cm]. This was not a surprise because the plants were extras, and thus were being held for an extended period whilst not receiving constant fertilization. Most nutrients were low: nitrogen (N) at 1.49 ppm, phosphorus (P) at 2.6 ppm, potassium (K) at 13.3 ppm, calcium (Ca) at 21.5 ppm, Mg at 6.71 ppm, and sulfur (S) at 29.1 ppm. With Mg being low, that supported the idea that a Mg deficiency was a possible cause.

Yet the substrate pH was 6.40. This was a little surprising because it negated the possibility that a low substrate pH induced micronutrient toxicity as being the cause, but on a positive note, it helped narrow down the diagnosis.

Leaf Tissue Test Results. The results from the leaf tissue analysis tests changed the initial diagnosis (all value interpretations are based on averages for floriculture species because no species-specific recommendations exist for Mexican sunflowers). While low N (2.44%), satisfactory P (0.20%), and satisfactory S (0.26%) concentrations were in the leaves, Mg was excessive (1.65%) and so was Ca (6.42%). In addition, K was considered deficient (1.06%) and Fe excessive at 1280 ppm. Mn (167 ppm), Zn (70.6 ppm), and Cu (7.59 ppm), were considered satisfactory and B was considered elevated at 63.3 ppm.



Figure 2. Initial symptoms of interveinal chlorosis on the lower foliage due to insufficient potassium (K). (Photo: Brian Whipker)



Figure 3. Lower leaf purplish-black coloration on Mexican sunflowers. On other species, low magnesium (Mg) can result in similar symptoms, but here potassium (K) was deficient and iron (Fe) was excessive in the leaf tissue. (Photo: Brian Whipker)

Mexican Sunflower

(*Tithonia rotundifolia*)




Figure 4. The progression of lower leaf interveinal chlorosis to necrosis in Mexican sunflowers due to low potassium and possibly excessive iron contributes to the purplish-black leaves. (Photo: Brian Whipker)

The high concentrations of Mg discount the possibility of a Mg deficiency. Excessive Ca concentrations can inhibit plant uptake of K and Mg. While K was low, it is doubtful that uptake was limited by the excessive levels of Ca because excessive Ca usually has a greater impact on Mg uptake and those levels were also excessive. This situation suggests that the lower leaf interveinal chlorosis was due to insufficient K fertilizer levels. This conclusion is further supported by the author's observations that plants with a higher proportion of stems usually require a higher supply of K. [Stem tissues usually require higher amounts of K and it is an important component of providing stem strength (Bryson et al., 2014)].

It is far easier to diagnose a problem when only one parameter is causing the issue. It is much more difficult to separate two factors that overlap. The excessive Fe concentrations at 1280 ppm cannot be ignored in the diagnosis. In general, Fe concentrations >800 ppm (Bryson et al., 2014) start to become problematic and leaf symptomology may appear due to toxic accumulations. The excessive Fe concentrations are the most likely what contributed to the lower leaf purplish-black discoloration, although

advanced K deficiencies will also result in total necrosis of the lower leaves. How the excessive Fe accumulated is still unknown because, with a substrate pH of 6.40, it is clear it was not caused by acidic substrate levels.

Conclusions.

The diagnosis is a combination of low K and excessive Fe as the most likely causes of this discoloration. But without clear-cut data to support this, one needs to acknowledge the limitations of this diagnosis. Conducting a K rate study to determine optimal K fertility levels and a substrate range experiment would help with clarity on this issue in Mexican sunflowers. In the meantime, consider providing higher levels of K fertilization to Mexican sunflowers and monitor the tissue for elevated Fe concentrations to hopefully avoid this issue.

Additional Resources:

Bryson, G.M. H.A. Mills, D.N. Sasseville, J.B. Jones, and A.V. Barker. 2014. Plant analysis handbook III. Micro-Macro Publ., Athens, GA.

e-GRO Alert

www.e-gro.org

CONTRIBUTORS

Dr. Nora Catlin
Floriculture Specialist
Cornell Cooperative Extension
Suffolk County
nora.catlin@cornell.edu

Dr. Chris Currey
Assistant Professor of Floriculture
Iowa State University
ccurrey@iastate.edu

Dr. Ryan Dickson
Greenhouse Horticulture and
Controlled-Environment Agriculture
University of Arkansas
rvand@uark.edu

Thomas Ford
Commercial Horticulture Educator
Penn State Extension
tf2@psu.edu

Dan Gilrein
Entomology Specialist
Cornell Cooperative Extension
Suffolk County
dog1@cornell.edu

Dr. Chieri Kubota
Controlled Environments Agriculture
The Ohio State University
kubota.10@osu.edu

Heidi Lindberg
Floriculture Extension Educator
Michigan State University
wollege@anr.msu.edu

Dr. Roberto Lopez
Floriculture Extension & Research
Michigan State University
relopez@msu.edu

Dr. Neil Mattson
Greenhouse Research & Extension
Cornell University
neil.mattson@cornell.edu

Dr. W. Garrett Owen
Sustainable Greenhouse & Nursery
Systems Extension & Research
The Ohio State University
owen.367@osu.edu

Dr. Rosa E. Raudales
Greenhouse Extension Specialist
University of Connecticut
rosa.raudales@uconn.edu

Dr. Alicia Rihn
Agricultural & Resource Economics
University of Tennessee-Knoxville
arihn@utk.edu

Dr. Debalina Saha
Horticulture Weed Science
Michigan State University
sahadeb2@msu.edu

Dr. Beth Scheckelhoff
Extension Educator - Greenhouse Systems
The Ohio State University
scheckelhoff.11@osu.edu

Dr. Ariana Torres-Bravo
Horticulture/ Ag. Economics
Purdue University
torres2@purdue.edu

Dr. Brian Whipker
Floriculture Extension & Research
NC State University
bwhipker@ncsu.edu

Dr. Jean Williams-Woodward
Ornamental Extension Plant Pathologist
University of Georgia
jwoodwar@uga.edu

Copyright © 2023

Where trade names, proprietary products, or specific equipment are listed, no discrimination is intended and no endorsement, guarantee or warranty is implied by the authors, universities or associations.

Cooperating Universities

Cornell CALS
College of Agriculture and Life Sciences

**Cornell Cooperative Extension
Suffolk County**



PennState Extension

IOWA STATE UNIVERSITY

**UTIA INSTITUTE OF
AGRICULTURE**
THE UNIVERSITY OF TENNESSEE

UCONN



**College of Agricultural &
Environmental Sciences**
UNIVERSITY OF GEORGIA

**MICHIGAN STATE
UNIVERSITY**

**U of A DIVISION OF AGRICULTURE
RESEARCH & EXTENSION**
University of Arkansas System

**PURDUE
UNIVERSITY**



**THE OHIO STATE
UNIVERSITY**

**NC STATE
UNIVERSITY**

In cooperation with our local and state greenhouse organizations

MAUMEE VALLEY GROWERS
Choose the Very Best.



Metro Detroit Flower Growers Association



**CONNECTICUT
GREENHOUSE
GROWERS
ASSOCIATION**



**Indiana
FLOWER
GROWERS
Association**

