

Some Viruses on Greenhouse Flower Crops:

Impatiens necrotic spot tospovirus (INSV)
Tomato spotted wilt tospovirus (TSWV)
Cucumber mosaic cucumovirus (CMV)
Tobacco mosaic tobamovirus (TMV)

Insects that may serve as virus vectors:

Aphids

Thrips

Whiteflies, Leafhoppers, Planthoppers

Beetles

Mealybugs

(and mites get in the act, too, at times!)

Flower break

Mosaic

Chlorotic Mottle

Ringspots

VIRUS SYMPTOMS:

Chlorotic or Necrotic Spots

Line Patterns

Leaf Strapping

Distortion

Stem cankers

Vein clearing, vein banding

Stunt

Flower Break Symptoms

TFBV
Tulip flower break virus



NGI Flower Break symptoms



Flower Break In Lily





Cucumber mosaic cucumovirus (CMV)

Angelonia Flower Break Virus Angelonia



Western Flower Thrips is an important vector of INSV and TSWV: Scout for thrips injury and you may find the virus symptoms too!



Scout for thrips!





- Injury and frass on leaves and petals
- YSC counts
- Breathe CO₂ into flowers

You can learn almost all the virus symptoms by looking at INSV and TSWV (they cause so many different symptoms on so many crops!)









Tool for Your Scouting: Immunostrips

Agdia Inc. agdia.com

Test whether a virus is the one you think . . .



INSV Symptoms

can appear on hundreds of different kinds of plant

Necrotic spots



New Guinea

INSV

Necrotic spots



Coleus - INSV




Symptoms of Impatiens Necrotic Spot Virus (INSV) on Pepper transplants


Necrotic spots

Chlorotic spots





Impatiens Necrotic Spot Virus





Trout Lily - INSV



Saintpaulia INSV



Cineraria - INSV



Leaf necrosis

Nemesia - INSV





Gloxinia - INSV




Tomato Spotted Wilt Virus

TSWV is Another orthotospovirus

(cousin to INSV)

To decide, use immunostrips



Verbena -TSWV

veinclearing



Zinnias -TSWV

Chlorosis, Vein clearing





Basil TSWV

Begonia - TSWV

Chlorotic mottle with 'green islands'







Senettis

Pericallis hyb.







NGI -TSWV

Necrotic stem cankers


Tomato Spotted Wilt Virus (TSWV)



mum



INSV on Rieger Begonia









Cucumber mosaic cucumovirus

Infects > 1000 spp.

Cheryl Smith, UNH Yellow mottle with ringspots

CMV on Monkshood

Insects that may serve as CMV vectors:

Aphids!
>80 species
Non-persistent
Stylet-borne



Black bean aphid Image by Sandra Jensen





Cheryl <u>cfrank@uvm.edu</u>



Questions?





Stephanie sburnett@maine.edu



Betsy eml38@cornell.edu



John jps3@cornell.edu



Elise ipmelise@gmail.com