

# Some Other Biocontrols



Natural Enemy ID  
Greenhouse Scout School  
**Session 7: March 5, 2026**

**Cheryl Frank Sullivan**  
University of Vermont  
Email: [cfrank@uvm.edu](mailto:cfrank@uvm.edu)



# Other Predators

## *Anystis baccarum*

### (Crazee Mite)

- Newly available (select states)
- Has erratic searching behavior 'whirligig mite'
- Generalist (attacks aphids, thrips, mites, and more, other biocontrol agents and each other)
- Need lots of prey and disperse well throughout crops.
- Research is ongoing



F. Piednoir; [iNaturalist](#)



I. Tuunainen; [iNaturalist](#)

# *Dicyphus hesperus*

- Predatory bug (as adults and nymphs)
- Attacks whiteflies, primarily.
- Will also feed on two-spotted spider mite, thrips and moth eggs.
- A bit slow to establish (often supported with mullein banker plants)
- Used primarily in vegetable production.
- Might cause plant damage in high numbers.




# Mullein Banker Plants



# Assassin Bugs

- Many species, Leafhopper Assassin Bug (*Zelus renardii*) commercially available.
- Generalist predators that will feed on leafhoppers, aphids, caterpillars, thrips, mites and more.
- Effective in many ecosystems (gardens, fields, orchards, greenhouses, high tunnels).
- Live ~2mo and eggs take about 10 days to hatch (they come as eggs).
- Predatory as nymphs and adults.



# Mantids

- Generalist predators that will feed on just about anything and themselves.
- Usually sold as egg cases
- Predatory as nymphs and adults.

Kai Squires, [iNaturalist](#)



Scot Nelson, [Flickr](#)



## Other Parasitic Wasps

### *Trichogramma* spp.

- Parasitize moth eggs
- Adults < 1 mm, larvae develop within entirely within moth eggs
- Parasitized eggs turn dark
- Several species commercially available for various moths and climates.



# *Diglyphus isaea*

- Parasitizes leaf miner larvae (warm weather releases)
- All stages but adults hidden within mines on foliage.
- Adults dark metallic blue/green but appears black; legs black/yellow; 1.5-2mm in size
- Adults also host feed.



# *Dacnusa sibirica*

- Parasitizes leaf miner larvae; prefers 1-2<sup>nd</sup> instars, (cool weather – spring/fall releases)
- Adults dark brown to black in color; 2 -3mm in size; long antennae; wings longer than abdomen.




# Other Pathogens

## Entomopathogenic fungi

- *Metarhizium brunneum* (thrips, whiteflies, thrips, aphids, mites and others). Does well for soil-based pests.
- *Isaria fumosorosea* (aphids, whiteflies, thrips, mites, flies and many more.)
- Spores need to contact host to be effective. They then penetrate exoskeleton, grow inside and release toxins killing pests, then emerge and sporulate to infect others.
- Other species can just show up.





# ***Bacillus thuringiensis*** **(*B.t.*)**

- Bacteria
- Produce spores & protein crystal toxins that kill pests
- Must be consumed to work
- Death occurs through starvation.
- Many strains available, primarily targets larvae (beetles, moths and flies).
  - B.t. var 'israelensis' (mosquitoes, black flies, & others)
  - B.t. var 'kurstaki' (many different caterpillars)

Aphid mummy created by a *Praon* sp., a wasp parasitoid



The Common Flower Fly, *Syrphus ribesii*, is a Syrphid.



Tobacco hornworm parasitized by wasp parasitoid (*Cotesia congregata*)



# To Know Them Is To Love Them!


The Swift Feather-legged Fly, *Trichopoda pennipes*, is a Tachinid.



Syrphid fly maggot attacking aphids.



# Thank You!



Cornell University



This work is supported by the National Institute of Food & Agric., U.S. Dept. of Agric., through the Northeast Sustainable Agric. Research & Education program under subaward number ONE22-419), UVM Extension and College of Agriculture and Life Sciences.

Any opinions, findings, conclusions, recommendations, use of tradenames and corporations expressed herein are those of the authors and do not necessarily reflect the view of or an endorsement by the sponsors or institutions. Images are subject to copyright and are used herein with permissions. For educational purposes only. Not for reproduction without permission from the author(s).